

Frequency and Etiology of Pediatric Trauma, Experience at Tertiary Care Hospital

Muhammad Kashif Bashir^{*1}, Aisha Ishtiaq², Shazia Bashir³

¹Department of Pediatric Surgery, King Edward Medical University/ Mayo Hospital Lahore, Pakistan.

²Department of Pediatric Surgery, Mayo Hospital Lahore, Lahore, Pakistan.

³Department of Trauma and Rehabilitation, Nasjon University of Health and Sciences, Lahore, Pakistan.

Abstract: Objective: To determine etiology and spectrum of trauma in children and to make recommendations for its prevention.

Materials and Methods: This Hospital based Retrospective observational study was conducted in Department of Pediatric Surgery King Edward Medical University / Mayo Hospital Lahore from March 2021 to February 2022. All patients presented in Pediatric Surgical emergency King Edward Medical University/ Mayo Hospital Lahore were included in this study. Data regarding age, gender, mechanism and severity of trauma and its management was analyzed and recorded on a prescribed Performa.

Results: During the study period a total of 3850 patients having trauma were presented in pediatric surgical emergency. There were 2206 (57.3%) male patients and 1644 (42.7%) females with male to female ratio of 1.3:1. Majority of children affected were 8 to 10 years age. Burn trauma was noted in 2400 (62.3%) while 1450 (37.7%) patients have poly trauma due to different etiological factors. Regarding burn trauma, majority of patients 1620 (67.5%) were having scald burn injury. Pedestrians 260 (17.9%) hitting with motor vehicle was noted to be major mechanism of trauma.

Conclusion: Motor vehicle collision is most common mechanism of pediatric trauma after pediatric burn injury. There is a need for parental education and strict implementation of traffic laws to prevent trauma at pediatric age group.

Keywords: Pediatric, Trauma, Etiology, Outcome, Management, Mortality.

INTRODUCTION

Trauma is recognized as major the cause of morbidity and mortality in children [1-3]. Trauma can either be blunt or penetrating depending on its etiology [4]. Road traffic accidents and burn injuries are major cause of trauma in children [5]. There is a need for vigilant assessment and management. According to one survey conducted at tertiary care hospital showed that road traffic accident is a major cause of mortality and morbidity in children. In Pakistan trauma is one of the most common causes of death in pediatric population [6-8]. Mayo hospital Lahore is one of the major tertiary care hospitals in country receiving all kinds of pediatric trauma. There is a need to know etiology, mechanism of injury, management and outcome of trauma in children. Therefore, this study was done to improve the care of such patients.

MATERIALS AND METHODS

This retrospective single centre study was done at department of pediatric surgical emergency King Edward Medical University/ Mayo hospital Lahore from March 2021 to February 2022. As it is a retrospective study so the hospital record admission files were the main source of information. Data was

collected including age, gender, clinical presentation at time of admission, geographical area of patient, type and mechanism of trauma. Clinical findings written on admission files were reviewed retrospectively and noted on a prescribed Performa and data was analyzed.

RESULTS

A total of 3850 patients having trauma with different etiology presented in department of pediatric surgical emergency. Regarding gender distribution there were 2206 (57.3%) male patients and 1644 (42.7%) females with male to female ratio of 1.3:1. Regarding age at time of presentation, there were 280 (7.3%) patients with age from 2 to 4 years, 800 (20.8%) patients with age of 5 to 7 years, 1850 (48.0%) patients with age range of 8 to 10 years and 920 (23.9%) patients were in age range of 11 to 12 years (Fig.1).




Fig. (1). Graph showing Age Range of Patients.

*Address correspondence to this author at the Department of Pediatric Surgery, King Edward Medical University/ Mayo Hospital Lahore, Pakistan.
Email: kashif.paeds.surgeon@gmail.com

Patients were divided into two groups. There were 2400 (n=1) patients with burn trauma and 1450 (n=2) patients having poly trauma due to different etiology. Regarding burn injuries, 1620 (67.5%) patients presented with scald burn injury, 265 (11%) with electric current, 480 (20%) with flame burn and 35 (1.5%) with chemical burn injuries (Table 1).

Table 1. Showing Pattern of Burn Trauma.

S.No	Pattern of burn trauma	No of patients (n=1)	Percentages
1	Scald burn	1620	67.5%
2	Flame burn	480	20%
3	Electric burn	265	11%
4	Chemical burn	35	1.5%
	Total	2400	100%

Regarding mechanism of injury in group 2 patients with poly trauma it was noted that majority of trauma patients were pedestrians having trauma by hitting with motor vehicles 260 (17.9%) followed by 665 (45.9%) Pedestrian hit by motor bike, 22 (2.8%) with history of fall. 150 (10.3%) children got trauma with bicycle handle bar injury and in 200 (13.8%) children due to sports trauma. trauma after gunshot injury was noted in 16 (1.1%) cases and in 4 (0.3) cases children got trauma after child abuse. 35 (2.4%) patients got trauma after dog bite while 80 (5.5%) patients have trauma due to different etiological factors (Table 2).

Table 2. Showing Different Mechanisms of Injury in Poly Trauma.

S. No	Mechanism of trauma	No of patients (n=2)	Percentages
1	Pedestrians hit by motor vehicle	260	17.9%
2	Pedestrians hit by motor bike	665	45.9%
3	Fire arm injury	16	1.1%
4	Bicycle handle bar injury	150	10.3%
5	Sports trauma	200	13.8%
6	History of fall	40	2.8%
7	Dog bite	35	2.4%
8	Child abuse	4	0.3%
9	Others	80	5.5%
	Total	1450	100%

DISCUSSION

It is seen in this study, that male children are more suffered from trauma than females. This is consistent with findings from other national and international studies which also

showed more involvement of male children [9-11]. Caught on to the fact that more children presented in month of June to August. This can be because of schools summer vacations and children have higher activity levels, greater freedom to play outside. Same results are also observed in another study [12,13]. In current study most commonly involved age group is from 5 to 10 years age. This can be due to the fact that at this age children have more chance to play outside their homes with their fellows where they can involved in more risky games. Similar findings also noted in another study [14]. Regarding mechanism of injury it is shown that more children were suffered from trauma after road traffic accident with motor vehicle or motor cycle followed by children injured by fall from roof or stairs. Toddlers and preschool children had also injury by having some heavy object fallen upon them. Similar results are also observed by other studies [15, 17]. In this study it was shown that abdomen and chest are most commonly involved areas in trauma. Same findings were noted in other studies [18, 20]. In abdominal trauma, we noted that liver and spleen were commonly injured organs. Same findings were also noted by other studies conducted at national and international level [21, 22]. In current study it is also noted that mostly children had liver and spleen injury of grade 2 to 3 except in few cases of severe poly trauma, where children had liver and spleen injury of more than 3 grade. Liver and spleen injury of grade 1 to 3 were managed conservative in our intensive care unit. In children with advance grade liver and spleen trauma exploratory laprotomy was performed. It is evident in other studies also that mostly liver and spleen trauma in children can be managed conservatively [23]. Among children suffered from chest trauma, pneumothorax and haemothorax were most commonly findings for which chest intubation was done. Similar results were also shown in other studies which showed that most commonly intervention needed in blunt chest trauma in children is chest intubation [24, 25].

CONCLUSION

Burn injuries and road traffic accidents are main cause of trauma in children leading to high mortality and morbidity. These injuries can be prevented by educating parents and children at electronic and social media. There should be special teaching classes for traffic laws and safety against home hazards. Strict implementation of traffic laws regarding safe speed and seat belt safety is recommended.

AUTHORS' CONTRIBUTION

Muhammad Kashif Bashir: Conception and design of the study, Data analysis, Manuscript writing.

Aisha Ishtiaq: Data Analysis, Critical review.

Shazia Bashir: Designing of the study, Data collection, Data entry and interpretation.

CONFLICT OF INTEREST

Declared none.

ACKNOWLEDGEMENTS

Declared none.

REFERENCES

- [1] Kant KC, Kant KS, Noman M. A study to evaluate the clinico-etiological profile and management of patients with trauma to the chest in a tertiary care Hospital. *Eur J Mol Clin Med* 2020; 7: 5975-82.
- [2] MdAnisuzzaman M, Hosain SN, Reza MM, Kibria MG, Ferdous S. Management of chest trauma in Bangladesh perspective: Experience of a decade. *Cardiovasc J* 2019; 12: 3-8.
- [3] Ibrahim SR, Abdelaziz AE. Retrospective statistical study of thoracic trauma patients in Al-hussein hospital, Al-azhar university. *Egypt J Hosp Med* 2021; 84: 1650-4.
- [4] Ludwig C, Koryllos A. Management of chest trauma. *J Thoracic Dis* 2017; 9: S172-7.
- [5] National Health Portal. World Trauma Day 2019. Available at: https://www.nhp.gov.in/world-trauma-day-2019_pg [updated 2019 Oct 21].
- [6] Mathangasinghe Y, Pradeep IHDS, Rasnayake D. Demographic, clinical features and outcome determinants of thoracic trauma in Sri Lanka: A multicentre prospective cohort study. *Canadian Respir J* 2020; Article ID 1219439.
- [7] Teisch LF, Allen CJ, Tashiro J, *et al.* Injury patterns and outcomes following pediatric bicycle accidents. *Pediatr Surg Int* 2015; 31: 1021-5.
- [8] Eberhardt CS, Zand T, Ceroni D, *et al.* The seatbelt syndrome – do we have a chance? A report of 3 cases with review of literature. *Pediatr Emerg Care* 2016; 32: 318-22.
- [9] Dai LN, Chen CD, Lin XK, *et al.* Abdominal injuries involving bicycle handlebars in 219 children: Results of 8 year follow-up. *Eur J Trauma Emerg Surg* 2015; 41: 551-5.
- [10] Choudhary A, Prasad K, Sreeramulu PN. A clinico—epidemiological study of traumatic chest injuries in a rural tertiary care centre in India: Our experience. *Int J Biomed Adv Res* 2015; 6: 110-4.
- [11] Mahajan P, Kuppermann N, Tunik M, *et al.* Comparison of clinician suspicion versus a clinical prediction rule in identifying children at risk for intra-abdominal injuries after blunt torso trauma. *Acad Emerg Med* 2015; 22: 1034-41.
- [12] Dandona R, Kumar GA, Gururaj G, *et al.* Mortality due to road injuries in the states of India: the global burden of disease study 1990–2017. *Lancet Publ Health* 2020; 5: e86-e98.
- [13] Djordjevic I, Slavkovic A, Marjanovic Z, Zivanovic D. Blunt trauma in paediatric patients - experience from a small centre. *West Indian Med J* 2015; 64(2): 126-30.
- [14] Notrica DM. Pediatric blunt abdominal trauma: Current management. *Curr Opin Crit Care* 2015; 21(6): 531-7.
- [15] LeeVan E, Zmora O, Cazzulino F, Burke RV, Zagory J, Upperman JS. Management of pediatric blunt renal trauma: A systematic review. *J Trauma Acute Care Surg* 2016; 80(3): 519-28.
- [16] Cunningham RM, Walton MA, Carter PM. The major causes of death in children and adolescents in the United States. *N Engl J Med* 2018; 379: 2468-75.
- [17] Kroczyk EK, Wieners G, Steffen I, *et al.* Non-traumatic incidental findings in patients undergoing whole-body computed tomography at initial emergency admission. *Emerg Med J* 2017; 34(10):643-6.
- [18] Linder F, Mani K, Juhlin C, Eklöf H. Routine whole body CT of high energy trauma patients leads to excessive radiation exposure. *Scand J Trauma Resusc Emerg Med* 2016; 24: 7
- [19] Botelho F, Truche P, Mooney DP, *et al.* Pediatric trauma primary survey performance among surgical and non-surgical pediatric providers in a Brazilian trauma center. *Trauma Surg Acute Care Open* 2020; 5: e000451.
- [20] Ahmad T, Haroon Khan M, Murad MA, *et al.* Research trends in rabies vaccine in the last three decades: A bibliometric analysis of global perspective. *Hum Vaccin Immunother* 2021; 17: 3169-77.
- [21] Karydakis P, Giakoumettis D, Themistocleous M. The 100 most cited papers about pediatric traumatic brain injury: A bibliometric analysis. *Ir J Med Sci* 2020; 189: 315-25.
- [22] Liu F, Wu TT, Lei G, *et al.* Worldwide tendency and perspectives in traumatic dental injuries: A bibliometric analysis over two decades (1999-2018). *Dent Traumatol* 2020; 36: 489-97.
- [23] Allahabadi S, Feeley SE, Lansdown DA, Pandya NK, Feeley BT. Influential articles on pediatric and adolescent anterior cruciate ligament injuries: A bibliometric analysis. *Orthop J Sports Med* 2021; 9: 1-9.
- [24] Tran BX, Pham TV, Ha GH, *et al.* A bibliometric analysis of the global research trend in child maltreatment. *Int J Environ Res Public Health* 2018; 15: 1456.
- [25] Brian JD, Ramanathan R, Francisco T, Robert LS, John G, Bjerke SH. Considerations in Pediatric Trauma. 2021; Available online at: <https://emedicine.medscape.com/article/435031-overview> [accessed July 16, 2021].